



The Oregon Department of Geology and Mineral Industries (DOGAMI) has been identifying and mapping the tsunami inundation hazard along the Oregon coast since 1994. In Oregon, DOGAMI manages the National Tsunami Hazard Mitigation Program, which has been administered by the National Oceanic and Atmospheric Administration (NOAA) since 1995. DOGAMI's work is designed to help cities, counties, and other sites in coastal areas reduce the potential for disastrous tsunami-related consequences by understanding and mitigating this geologic hazard. Using federal funding awarded by NOAA, DOGAMI has developed a new generation of tsunami inundation maps to help residents and visitors along the entire Oregon coast prepare for the next Cascadia Subduction Zone (CSZ) earthquake and tsunami.

The CSZ is the tectonic plate boundary between the North American Plate and the Juan de Fuca Plate (Figure 1). These plates are converging at a rate of about 1.5 inches per year, but the movement is not smooth and continuous. Rather, the plates lock in place, and unreleased energy builds over time. At intervals, this accumulated energy is violently released in the form of a megathrust earthquake rupture, where the North American Plate suddenly slips westward over the Juan de Fuca Plate. This rupture causes a vertical displacement of water that creates a tsunami (Figure 2). Similar rupture processes and tsunamis have occurred elsewhere on the planet where subduction zones exist: for example, offshore Chile in 1960 and 2010, offshore Alaska in 1964, near Sumatra in 2004, and offshore Japan in March 2011.

CSZ frequency: Comprehensive research of the offshore geologic record indicates that at least 19 major ruptures of the full length of the CSZ have occurred off the Oregon coast over the past 10,000 years (Figure 3). All 19 of these full-rupture CSZ events were likely magnitude 8.9 to 9.2 earthquakes (Witter and others, 2011). The most recent CSZ event happened approximately 300 years ago on January 26, 1700. Sand deposits carried onshore and left by the 1700 event have been found 12 miles inland, older tsunami sand deposits have also been discovered in estuaries 6 miles inland. As shown in Figure 3, the range in time between these 19 events varies from 110 to 1,150 years, with a median time interval of 490 years. In 2008 the United States Geological Survey (USGS) released the results of a study announcing that the probability of a magnitude 8.9 CSZ earthquake occurring over the next 30 years is 10% and that such earthquakes occur about every 500 years (WGCEP 2008).

CS2 Model Specifications: The sizes of the earthquake and its resultant tsunami are primarily driven by the amount and geometry of the slip that takes place when the North American Plate slips westward over the Juan de Fuca Plate during a CS2 event. DOGAMI has modeled a wide range of earthquake and tsunami sizes that take into account different fault geometries that could amplify the amount of seawater displacement and increase tsunami inundation. Seismic geophysical profiles show that there may be a steep splay fault running nearly parallel to the CS2 but closer to the Oregon coastline (Figure 1). The effect of this splay fault moving during a full-rupture CS2 event would be an increase in the amount of vertical displacement of the Pacific Ocean, resulting in an increase of the tsunami inundation onshore in

A cross-sectional diagram of the Juan de Fuca Plate subducting beneath the North American Plate. The Pacific Plate is shown to the west, with the Juan de Fuca Plate extending from it. The Juan de Fuca Plate is shown dipping into the mantle beneath the North American Plate. The subducting plate is labeled 'Juan de Fuca Plate' and the overriding plate is labeled 'NORTH AMERICAN PLATE'. The boundary between them is labeled 'JUAN DE FUCA RIFT ZONE'. The subducting plate is shown with a 'Locked Zone' and 'Megaseismic' areas. The overriding plate is shown with 'BIOLOGICAL DIVERSITY' and 'WATERSHED' features. The diagram illustrates the process of plate tectonics and the resulting geological features.

Figure 1: This block diagram depicts the tectonic setting of the region. See Figure 2 for the sequence of events that occur during a Cascadia Subduction Zone megathrust earthquake and tsunami.

A

B

C

D

E

Figure 2: The North American Plate rises over the descending Juan de Fuca Plate at a rate of approximately 1.5 inches per year.

Because the two plates are stuck in place at the "locked zone," ocean bulges up over time, and the North American Plate bulges up.

Eventually the locked zone ruptures and causes a great earthquake. The sudden slip of the two plates displaces Pacific Ocean water upward and creates a tsunami.

Displaced and uplifted Pacific Ocean water travels in all directions.

Along the Oregon coast, tsunami waves run up onto the land for several hours.

Figure 3 is a chart depicting the timing, frequency, and magnitude of the last 19 great Cascadia Subduction Zone events over the past 10,000 years. The x-axis represents the Research-Indicated Radiocarbon Age of C2 Event (Most recent in January 2002), ranging from 0 to 10,000 years. The y-axis represents the Number of Earthquakes and the Size of the Tsunami, ranging from 0 to 19. The chart shows 19 events, with the largest event (M9.0) occurring around 1700 years ago, and the smallest event (M7.0) occurring around 100 years ago. The chart also shows the timing of the 1997-1998 El Niño event and the 1992-1993 El Niño event. The chart is divided into two sections: 'larger but much less frequent tsunamis' (top) and 'smaller but more frequent tsunamis' (bottom). The chart is titled 'Figure 3: This chart depicts the timing, frequency, and magnitude of the last 19 great Cascadia Subduction Zone events over the past 10,000 years. The most recent event occurred on January 26, 2002. The 1700 event is considered to be a medium-sized event. The data used to create this chart came from research that examined the many submarine landslides known as 'narragansets' that are triggered only by these great earthquakes (Witter and others, 2011). The close correlation is "the bigger the landslide, the bigger the earthquake."

Table 1: Percent of Buildings within Tsunami Zones

	Entire Map Area	Uncorporated Areas
Total Buildings	24	24
Buildings within Tsunami Zones*		
Small	0	0
Medium	0	0
Large	0	0
Extra Large	0	0
Extra Extra Large	0	0

Figure 1: Percent of Buildings within Tsunami Zones

	Entire Map Area	Uncorporated Areas
Small	0.0%	0.0%
Medium	0.0%	0.0%
Large	0.0%	0.0%
Extra Large	0.0%	0.0%
Extra Extra Large	0.0%	0.0%

*Building counts shown are based on polygon centroids and are cumulative within the map area.

Figure 2: Cumulative Percent of Buildings within Tsunami Zones for Entire Map Area

Category	Percent of Buildings within Tsunami Zones
Small	0%
Medium	0%
Large	0%
Extra Large	0%
Extra Extra Large	0%

Figure 4: The table and chart show the number of buildings inundated for each "tsunami T-shirt scenario" for cities and unincorporated portions of the map.

A HUMBUG MOUNTAIN - NORTH A'

Elevation, feet (NAVD83)

Distance along Profile, miles

B HUMBUG MOUNTAIN - SOUTH B'

Elevation, feet (NAVD83)

Distance along Profile, miles

4079' line

46172' line

Pacific Ocean

XL

XI

L

S

Tsumami Scenarios

Figure 6: These profiles depict the expected maximum tsunami wave elevation for the five "tsunami T-shirt scenarios" along lines A-A' and B-B'. The tsunami scenarios are modeled to occur at high tide and to account for local subsidence or uplift of the ground surface.

Oregon. DOWAMI has also incorporated physical evidence that suggests that portions of the coast may drop 4 to 10 feet during the earthquake; this effect is known as subsidence. Detailed information on fault geometries, subsidence, computer models, and the methodology used to create the tsunami scenarios presented on this map can be found in DOWAMI Special Papers 41 (Priest and others, 2009) and 43 (Witter and others, 2011).

This tsunami inundation map displays the output of computer models representing five selected tsunami scenarios, all of which include the earthquake-produced subsidence and the tsunami-amplifying effects of the spillof fault. Each scenario assumes that a tsunami occurs at Mean Higher High Water (MHHW) tide; MHHW is defined as the average height of the higher high tides observed over an 18-year period at the Port of Long Beach. To make it easier to understand this scientific material and to enhance the educational aspects of hazard mitigation and response, the five scenarios are labeled as "T-shirt sizes" ranging from Small, Medium, Large, Extra Large, to Extra Extra Large (S, M, L, XL, XXL). The map legend depicts the respective amounts of spill, the frequency of occurrence, and the earthquake magnitude for these five scenarios. Figure 4 shows the cumulative number of buildings inundated within the map area.

The computer simulation model output is provided to DOGAMI as millions of points with values that indicate whether the location of each point is wet or dry. These points are converted to wet and dry contour lines that form the extent of inundation. The transition area between the wet and dry contour lines is termed the Wet/Dry Zone, which equates to the amount of error in the model when determining the maximum inundation for each scenario. Only the XXL Wet/Dry Zone is shown on this map.

This map also shows the regulatory tsunami inundation line (Oregon Revised Statutes 455.446 and 455.447), commonly known as the Senate Bill 379 line. Senate Bill 379 (1995) instructed DOGAMI to establish the area of expected tsunami inundation based on scientific evidence and tsunami modeling in order to prohibit the construction of new essential and special occupancy structures in this tsunami inundation zone (Priest, 1995).

Time Series Graphs and Wave Elevation Profiles In addition to the time series scenarios, the computer model produces time series data for "gauge" locations in the area. These points are simulated gauge stations that record the time, in seconds, of the tsunami wave arrival and the wave height observed. It is especially noteworthy that the greatest wave height and velocity observed are not necessarily associated with the first tsunami wave to arrive onshore. Therefore, evacuees should not assume that the tsunami event is over until the proper authorities have sounded the all-clear signal at the end of the evacuation. Figure 5 depicts the tsunami waves as they arrive at a simulated gauge station. Figure 6 depicts the overall wave height and inundation extent for all five scenarios at the profile locations shown on this map.

2012



Earthquake Size	Average Slip Range (ft)	Maximum Slip Range (ft)	Time to Accumulate Slip (yrs)	Earthquake Magnitude
XXL	59 to 72	118 to 144	1,200	-9.1
XL	56 to 72	115 to 144	1,050 to 1,200	-9.1
L	36 to 49	72 to 98	650 to 800	-9.0
M	23 to 30	46 to 62	425 to 525	-8.9
S	13 to 16	30 to 36	300	-8.7
XXL Wet/Dry Zone				

	Urban Growth Boundary		Fire Station
	Building Footprint		Police Station
	Simulated Gauge Station		School
	Profile Location		Hospital/Urgent Care Clinic
	Senate Bill 379 Line		US Highway
	State Park		State Highway
	Elevation Contour (25 ft intervals up to 200 ft)		Improved Road

Curr-01 Langlois	Curr-09 Gold Beach
Curr-02 Cape Blanco	Curr-10 Cape Sebastian
Curr-03 Denmark	Curr-11 Pistol River
Curr-04 Glenbrook	Curr-12 Carpenterville
Curr-05 Humboldt Mountain	Curr-13 Harris Beach
Curr-06 Sisters Fork	Curr-14 Orino River
Curr-07 Nesika Bay	Curr-15 Winchuck River
Curr-08 Northogue River	Curr-16 Brookings

Source Data:
This map is based on hydrodynamic tsunami modeling by Joseph Zhang, Oregon Health and Science University, Portland, Oregon. Model data were created by John T. English and George R. Priest, Department of Geology and Mineral Industries (DOGAMI), Portland, Oregon.
Hydrology data, contours, critical facilities, and building footprints were created by DOGAMI. Seane B119 379 data were redigitized by Rachel R. Lyles Smith and Sean G. Pickner, DOGAMI, in 2011 (GIS file set, in press, 2012).
Urban growth boundaries (2010) were provided by the Oregon Department of Land Conservation and Development (DLCD).
Transportation data (2010) provided by Curry County were edited by DOGAMI to improve the spatial accuracy of the features or to add newly constructed roads not present in the original data layer.
Lidar data from DOGAMI Lidar Data Quadrangle LDO-2009-42124-4-4-Portland.

[illegible]

Software: Esri ArcGIS® 10.0, Microsoft® Excel®, and Adobe® Illustrator®

Funding: This map was funded under award #NA09NW5467004 by the National Oceanic and Atmospheric Administration (NOAA) through the National Tsunami Hazard Mitigation Program.

Map Data Creation/Development:
Research Foundation Scanner: George R. Priest, Laura L. Stimmel, Daniel E. Cox, Paul A. Ferro, Sean G. Pickner, Rachel R. Lyles Smith
Research Data: Kaleena L.B. Hughes, Sean G. Pickner

Map Production:
Cartography: Kaleena L.B. Hughes, Sean G. Pickner, Taylore E. Womble
Text: Don W. Lewis, Rachel R. Lyles Smith
Editing: Don W. Lewis, Rachel R. Lyles Smith
Publication: Deborah A. Schuster

Map Date: 09/13/2012

