



The Oregon Department of Geology and Mineral Industries (DOGAMI) has been identifying and mapping the tsunami inundation hazard along the Oregon coast since 1994. In Oregon, DOGAMI manages the National Tsunami Hazard Mitigation Program, which has been administered by the National Oceanic and Atmospheric Administration (NOAA) since 1995. DOGAMI's work is designed to help cities, counties, and other states in coastal areas reduce the potential for disastrous tsunami-related consequences by understanding and mitigating this geologic hazard. Using federal funding awarded by NOAA, DOGAMI has developed a new generation of tsunami inundation maps to help residents and visitors along the entire Oregon coast prepare for the next Cascadia Subduction Zone (CSZ) earthquake and tsunami.

The CSZ is the tectonic plate boundary between the North American Plate and the Juan de Fuca Plate (Figure 1). These plates are converging at a rate of about 1.5 inches per year, but the movement is not smooth and continuous. Rather, the plates lock in place, and unreleased energy builds over time. At intervals, this accumulated energy is violently released in the form of a megathrust earthquake rupture, where the North American Plate suddenly slips westward over the Juan de Fuca Plate. This rupture causes a vertical displacement of water that creates a tsunami (Figure 2). Similar rupture processes and tsunamis have occurred elsewhere on the planet where subduction zones exist: for example, offshore Chile in 1960 and 2010, offshore Alaska in 1964, near Sumatra in 2004, and offshores Japan in March 2011.

Oregon, DOGAMI has also incorporated physical evidence that suggests that portions of the coast may drop 4 to 10 feet during the earthquake; this effect is known as subsidence. Detailed information on fault geometries, subsidence, computer models, and the methodology used to create the tsunami scenarios presented on this map can be found in DOGAMI Special Papers 41 (Priest and others, 2009) and 43 (Witter and others, 2011).

This tsunami inundation map displays the output of computer models representing five selected tsunami scenarios, all of which include the earthquake-produced subsidence and the tsunami-amplifying effects of the geyser fault. Each scenario assumes that a tsunami occurs at Mean Higher High Water (MHHW) tide; MHHW is defined as the average height of the highest high tides observed over a 18-yr period at the Yaquina Bay (Central Coast Model) tide gauge. To make it easier to understand this scientific material and to enhance the educational aspects of hazard mitigation and response, the five scenarios are labeled as "T-shirt sizes" ranging from Small, Medium, Large, Extra Large, to Extra Extra Large (S, M, L, XL, XXL). The map legend depicts the respective amounts of slip, the frequency of occurrence, and the earthquake magnitude for these five scenarios. Figure 4 shows the cumulative number of buildings inundated within the map area.

The computer simulation model output is provided to DQAGMI as millions of points with values that indicate whether the location of each point is wet or dry. These points are converted to wet and dry contour lines that form the extent of inundation. The transition area between the wet and dry contour lines is termed the Wet/Dry Zone, which equates to the amount of error in the model when determining the maximum inundation for each scenario. Only the XXI Wet/Dry Zone is shown on this map.

CSZ Model Specifications: The sizes of the earthquake and its resultant tsunami are primarily driven by the amount and geometry of the slip at the plate interface. When the North American Plate slips westward over the Juan de Fuca Plate during a CSZ event, DOGAMI has modeled a wide range of earthquake and tsunami sizes that take into account different plate geometries that could amplify the amount of seawater displacement and increase tsunami inundation. Seismic geophysical profiles show that there may be a steep slip fault running nearly parallel to the CSZ but closer to the Oregon coastline (Figure 1). The effect of this slip fault moving during a full-rupture CSZ event would be an increase in the amount of vertical displacement of the Pacific Ocean, resulting in an increase of the tsunami inundation onshore in

A cross-sectional diagram of the Juan de Fuca Plate subducting beneath the North American Plate. The Pacific Plate is shown on the left, with the Juan de Fuca Plate extending from it. The Juan de Fuca Plate is shown subducting under the North American Plate. The subducting plate is labeled 'Juan de Fuca Plate' and the overriding plate is labeled 'NORTH AMERICAN PLATE'. The boundary between them is labeled 'Juan de Fuca Plate' and 'Locked Zone'. The subducting plate is shown dipping into the mantle, with a 'Magma' source indicated. The overriding plate is shown with a 'Magma' source. The diagram also shows the 'Cascades' and 'Willamette' regions.

Figure 1: This block diagram depicts the tectonic setting of the region. See Figure 2 for the sequence of events that occur during a Cascadia Subduction Zone megathrust earthquake and tsunami.

Figure 2: The North American Plate rides over the descending Juan de Fuca Plate at a rate of approximately 1.5 inches per year.

A The diagram shows the Pacific Plate subducting beneath the North American Plate. The Juan de Fuca Plate is labeled, and the North American Plate is shown moving over it.

B The plates are stuck in place at the 'locked zone'. Strain builds up over time, and the North American Plate begins to bulge up.

C The locked zone ruptures, releasing energy as a megathrust earthquake. The Pacific Plate is displaced northward, creating a tsunami.

D The displaced and uplifted Pacific Ocean water rushes in all directions, creating tsunami waves that travel for several hours.

	Entire Map Area	Unincorporated Areas
Total Buildings	169	169
Buildings within Tsunami Zones*		
Small	2	2
Medium	5	5
Large	8	8
Extra Large	19	19
Extra Extra Large	19	19
Percent of Buildings within Tsunami Zones		
Small	1.2%	1.2%
Medium	3.0%	3.0%
Large	4.7%	4.7%
Extra Large	11.2%	11.2%
Extra Extra Large	11.2%	11.2%

*Building counts shown are based on polygon centroids and are cumulative within the map area.

Category	Bar 1 (Approx. %)	Bar 2 (Approx. %)
Entire Map Area	5	10
Unincorporated Areas	5	10

Figure 4: The table and chart show the number of buildings inundated for each "tsunami T-shirt scenario" for cities

2013





Figure 6: These profiles depict the expected maximum tsunami wave elevation for the five "tsunami T-shirt scenarios" along lines A-A' and B-B'. The tsunami scenarios are modeled to occur at high tide and to account for local subsidence or uplift of the ground surface.



01

02

03

04

05

06

07

08

09

Douglas
Coco

OREGON

Douglas 01 Silver Lake
Douglas 02 Tahkenneth Lake
Douglas 03 Garden Lake
Douglas 04 Redport
Douglas 05 Bearport
Douglas 06 Douglas
Douglas 07 Sulphur Springs
Douglas 08 Umpqua River East
Douglas 09 Clear Lake

[illegible]